



## NETWORK 7 GUIDELINES FOR CLINICAL OUTCOMES

### ADULT CONVENTIONAL HEMODIALYSIS & PERITONEAL DIALYSIS

Clinical Indicator	Network 7 Goal	Notes
<b>Adequacy</b>		
Kt/V $\geq$ 1.2 - HD $\geq$ 1.7 - PD	96.1% 94.0%	Percentage of patients having Kt/V $\Rightarrow$ 1.2 Percentage of PD patients having Kt/V $\Rightarrow$ 1.7
URR $\geq$ 65%	95.5%	Percentage of patients having URR $\Rightarrow$ 65%
<b>Anemia</b>		
HGB <10g/dL	3.8%	Percentage of patients having Hgb less than 10
% Saturation $\Rightarrow$ 20	84.9%	Percentage of patients having %TSAT equal or greater than 20
Ferritin $\Rightarrow$ 100 & $\Rightarrow$ 1000ng/ml	85.6%	Percentage of patients having Ferritin equal or greater than 100 & equal or less than 1000
<b>Nutrition &amp; Bone Mineral Metabolism</b>		
Albumin (BCG/BCP) > 4.0/3.7g/dL	39.1%	Percentage of patients having Albumin greater than 4.0 / 3.7
Albumin (BCG/BCP) >3.5/3.2g/dL	86.0%	Percentage of patients having Albumin greater than 3.5 / 3.2
Calcium $\Rightarrow$ <10.2mg/dL	97.3%	Percentage of patients having Calcium equal or less than 10.2
Calcium $\Rightarrow$ <9.5mg/dL	73.0%	Percentage of patients having Calcium equal or less than 9.5
Phosphorus $\Rightarrow$ <5.5mg/dL	61.0%	Percentage of patients having PO4 equal or less than 5.5
<b>Vascular Access</b>		
AVF	66%	Percentage of patients having AVF in use
Catheter only $\geq$ 90 days	$\Rightarrow$ <10%	Percentage of patients with CVC only equal or greater than 90days